

## PRELIMINARY REPORT

CLIENT DETAILS -

LABORATORY DETAILS -

Contact

John Fox

Client

NAMOI COTTON LIMITED

Address PO BOX 1333

TOOWOOMBA QLD 4350

Manager

Laboratory

Address

SGS Melbourne EH&S

Adam Atkinson

10/585 Blackburn Road

Notting Hill Victoria 3168

Telephone

0429 903 079

Facsimile

61 7 46316184 jfox@namoicotton.com.au

Email

Samples

Facsimile Email

Telephone

+61395743200 +61395743399

Au.SampleReceipt.Melbourne@sgs.com

Project Order Number Merah North Cotton Gin Storm water Disch TBA 2

SGS Reference

Date Received Date Reported ME319267 R0 09 Mar 2021

COMMENTS

This is an interim report. Final QC checks are yet to be completed.

TKN & TP analysis subcontracted to SGS Sydney, Unit 16 33 Maddox St Alexandria NSW 2015, NATA Accreditation Number: 2562, Site Number: 4354, <ENTER REPORT NUMBER>.

SIGNATORIES

Christopher BENNETT Laboratory Technician

Ryan ZHANG

Inorganics Team Leader

Vanessa PALAMARA Senior Chemist

SGS Australia Pty Ltd ABN 44 000 964 278

Environment, Health and

Bldg 10, 585 Blackburn Rd

Notting Hill VIC

t +61 3 9574 3200 f +61 3 9574 3399

www.sgs.com.au



2-fluorobiphenyl (Surrogate)

## PRELIMINARY REPORT

ME319267 R0

	Sample Number Sample Matrix Sample Date Sample Name			ME319267.002 Water 25/2/21 14:00 Point19	
Parameter	Units	LOR			
Total and Volatile Suspended Solids (TSS / VSS) Me	thod: AN114 Teste	ed: 15/3/2021			
Total Suspended Solids Dried at 103-105°C	mg/L	5	610	460	
Nitrate Nitrogen and Nitrite Nitrogen (NOx) by FIA M	ethod: MA-1127-04	Tested: 12/3	8/2021		
Nitrate/Nitrite Nitrogen, NOx as N	mg/L	0.01	0.38	0.57	
TKN Kjeldahl Digestion by Discrete Analyser Method  Total Kjeldahl Nitrogen	d: AN292 Tested:	0.05	NVL	NVL	
Total Nitrogen (calc)	mg/L	0.05	NVL	NVL	
Total Phosphorus by Kjeldahl Digestion DA in Water	Method: AN279/AN				
Total Phosphorus (Kjeldahl Digestion) as P	mg/L	0.02	NVL	NVL	
SVOC Compounds in Waters by GC-QQQ Method: N 4,4-DDD	μg/L	1	<1	<1	
4,4-DDE	µg/L	1	<1	<1	
4,4-DDT	µg/L	1	<1	<1	
o,p-DDT	μg/L	1	<1	<1	
Aldrin	μg/L	1	<1	<1	
alpha-BHC	μg/L	1	<1	<1	
alpha-Chlordane	μg/L	1	<1	<1	
beta-BHC Chlorpyrifos	μg/L	1	<1 <1	<1 <1	
delta-BHC	μg/L μg/L	1	<1	<1	
Dieldrin	μg/L	1	<1	<1	
Dimethoate	µg/L	1	<1	<1	
Endosulfan 1	μg/L	1	<1	<1	
Endosulfan 2	µg/L	1	<1	<1	
Endosulfan Sulphate	μg/L	1	<1	<1	
Endrin	μg/L	1	<1	<1	
Endrin Aldehyde	μg/L	1	<1	<1	
Endrin ketone	μg/L	1	<1	<1	
gamma-BHC	μg/L	1	<1	<1	
gamma-Chlordane	μg/L	1	<1	<1	
Heptachlor Epoxide	μg/L	1	<1	<1	
Hexachlorobenzene	μg/L	1	<1	<1	
Isodrin	μg/L	1	<1	<1	
Methoxychlor	μg/L	1	<1	<1	
Mirex	μg/L	1	<1	<1	

16-March-2021 Page 2 of 7



## PRELIMINARY REPORT

ME319267 R0

	•	Sample Number Sample Matrix Sample Date Sample Name	ME319267.001 Water 25/2/21 14:00 Point18	ME319267.002 Water 25/2/21 14:00 Point19
Parameter	Units	LOR		
Combined SVOC Pesticides in Water Method: MA 827	0 Tested: 11/3/20	021		
Dichlorvos	μg/L	1	<1	<1
Diazinon*	μg/L	1	<1	<1
Fenitrothion	μg/L	1	<1	<1
Malathion	μg/L	1	<1	<1
Parathion	μg/L	1	<1	<1
Bromophos ethyl*	μg/L	1	<1	<1
Methidathion	μg/L	1	<1	<1
Ethion	μg/L	1	<1	<1
Azinphos-methyl	μg/L	1	<1	<1
Surrogates		'		
d14-p-terphenyl (Surrogate)	%	-	-	-

16-March-2021 Page 3 of 7



## **QC SUMMARY**

MB blank results are compared to the Limit of Reporting

LCS and MS spike recoveries are measured as the percentage of analyte recovered from the sample compared the the amount of analyte spiked into the sample.

DUP and MSD relative percent differences are measured against their original counterpart samples according to the formula: the absolute difference of the two results divided by the average of the two results as a percentage. Where the DUP RPD is 'NA', the results are less than the LOR and thus the RPD is not applicable.

#### Combined SVOC Pesticides in Water Method: MA 8270

Parameter	QC Reference	Units	LOR	MB	LCS %Recovery
Dichlorvos	LB039698	μg/L	1	<1	123%
Diazinon	LB039698	μg/L	1	<1	NA
Fenitrothion	LB039698	μg/L	1	<1	NA
Malathion	LB039698	μg/L	1	<1	NA
Parathion	LB039698	μg/L	1	<1	NA
Bromophos ethyl	LB039698	μg/L	1	<1	NA
Methidathion	LB039698	μg/L	1	<1	NA
Ethion	LB039698	μg/L	1	<1	NA
Azinphos-methyl	LB039698	μg/L	1	<1	NA

#### Nitrate Nitrogen and Nitrite Nitrogen (NOx) by FIA Method: MA-1127-04

	Parameter	QC	Units	LOR	MB	DUP %RPD	LCS	MS	MSD %RPD
		Reference					%Recovery	%Recovery	
ı	Nitrate/Nitrite Nitrogen, NOx as N	LB039724	mg/L	0.01	<0.01	1%	NA	NA	NA

#### SVOC Compounds in Waters by GC-QQQ Method: MA8270

Parameter	QC Reference	Units	LOR	МВ	LCS %Recovery	
4,4-DDD	LB039698	μg/L	1	<1	NA	
4,4-DDE	LB039698	μg/L	1	<1	NA	
4,4-DDT	LB039698	μg/L	1	<1		
o,p-DDT	LB039698	μg/L	1	<1	NA	
Aldrin	LB039698	μg/L	1	<1	76%	
alpha-BHC	LB039698	μg/L	1	<1	NA	
alpha-Chlordane	LB039698	μg/L	1	<1	NA	
beta-BHC	LB039698	μg/L	1	<1	NA	
Chlorpyrifos	LB039698	μg/L	1	<1	NA	
delta-BHC	LB039698	μg/L	1	<1	NA	
Dieldrin	LB039698	μg/L	1	<1	69%	
Dimethoate	LB039698	μg/L	1	<1	NA	
Endosulfan 1	LB039698	μg/L	1	<1	NA	
Endosulfan 2	LB039698	μg/L	1	<1	NA	
Endosulfan Sulphate	LB039698	μg/L	1	<1	NA	
Endrin	LB039698	μg/L	1	<1	89%	
Endrin Aldehyde	LB039698	μg/L	1	<1	NA	
Endrin ketone	LB039698	μg/L	1	<1	NA	
gamma-BHC	LB039698	μg/L	1	<1	74%	
gamma-Chlordane	LB039698	μg/L	1	<1	NA	
Heptachlor Epoxide	LB039698	μg/L	1	<1	NA	
Hexachlorobenzene	LB039698	μg/L	1	<1	NA	
Isodrin	LB039698	μg/L	1	<1	NA	
Methoxychlor	LB039698	μg/L	1	<1	NA	
Mirex	LB039698	μg/L	1	<1	NA	
2-fluorobiphenyl (Surrogate)	LB039698	%	-	87%	70%	

16-March-2021 Page 4 of 7



## **QC SUMMARY**



MB blank results are compared to the Limit of Reporting

LCS and MS spike recoveries are measured as the percentage of analyte recovered from the sample compared the the amount of analyte spiked into the sample.

DUP and MSD relative percent differences are measured against their original counterpart samples according to the formula: the absolute difference of the two results divided by the average of the two results as a percentage. Where the DUP RPD is 'NA', the results are less than the LOR and thus the RPD is not applicable.

#### Total and Volatile Suspended Solids (TSS / VSS) Method: ME-(AU)-[ENV]AN114

Parameter	QC	Units	LOR	MB	DUP %RPD	LCS
	Reference					%Recovery
Total Suspended Solids Dried at 103-105°C	LB039753	mg/L	5	<5	11%	109%

16-March-2021 Page 5 of 7



## METHOD SUMMARY



METHOD

METHODOLOGY SUMMARY

AN114

Total Suspended and Volatile Suspended Solids: The sample is homogenised by shaking and a known volume is filtered through a pre-weighed GF/C filter paper and washed well with deionised water. The filter paper is dried and reweighed. The TSS is the residue retained by the filter per unit volume of sample. Reference APHA 2540 D. Internal Reference AN114

AN279/AN293(Sydney)

The sample is digested with Sulphuric acid, K2SO4 and CuSO4. All forms of phosphorus are converted into orthophosphate. The digest is cooled and placed on the discrete analyser for colorimetric analysis.

AN281

An unfiltered water or soil sample is first digested in a block digestor with sulfuric acid, K2SO4 and CuSO4. The ammonia produced following digestion is then measured colourimetrically using the Aquakem 250 Discrete Analyser. A portion of the digested sample is buffered to an alkaline pH, and interfering cations are complexed. The ammonia then reacts with salicylate and hypochlorite to give a blue colour whose absorbance is measured at 660nm and compared with calibration standards. This is proportional to the concentration of Total Kjeldahl Nitrogen in the original sample.

MA1127-04

Determination of nitrate/nitrite by Flow Injection Analysis (FIA). The method is based on reactions that are specific for the nitrite (NO2-) ion. Nitrate is quantitatively reduced to nitrite by passage of the sample through a copperised cadmium column. The nitrite (reduced nitrate plus original nitrite) is then determined by diazotisation with sulphanilamide under acidic conditions to form a diazonium ion. The diazonium ion is then coupled with N-(1-naphthyl)ethylenediamine dihydrochloride. The resulting pink dye absorbs at 540 nm. Nitrate concentrations are obtained by subtracting nitrite values, which have been previously analysed

MA8270

This method covers analytical procedures for the analysis of semi-volatile organic compounds (SVOC) including most neutral, acidic, and basic organic compounds based on the USEPA method 8270D. Samples are extracted into a solvent appropriate to the matrix and analysed using a gas chromatograph – triple quadrapole (GC–QQQ).

MS-EN-MA8270 Pest

This method covers analytical procedures for the analysis of semi-volatile organic compounds (SVOC pesticides) including most neutral, acidic, and basic organic compounds based on the USEPA method 8270D. Samples are extracted into a solvent appropriate to the matrix and analysed using a gas chromatograph – triple quadrapole (GC–QQQ).

16-March-2021 Page 6 of 7



# FOOTNOTES



FOOTNOTES

IS Insufficient sample for analysis. LOR Limit of Reporting LNR Sample listed, but not received. Raised or Lowered Limit of Reporting ↑↓ NATA accreditation does not cover the OFH QC result is above the upper tolerance performance of this service QFL QC result is below the lower tolerance Indicative data, theoretical holding time exceeded. The sample was not analysed for this analyte Indicates that both \* and \*\* apply. NVI Not Validated

Unless it is reported that sampling has been performed by SGS, the samples have been analysed as received.

Where "Total" analyte groups are reported (for example, Total PAHs, Total OC Pesticides) the total will be calculated as the sum of the individual analytes, with those analytes that are reported as <LOR being assumed to be zero. The summed (Total) limit of reporting is calculated by summing the individual analyte LORs and dividing by two. For example, where 16 individual analytes are being summed and each has an LOR of 0.1 mg/kg, the "Totals" LOR will be 1.6 / 2 (0.8 mg/kg). Where only 2 analytes are being summed, the "Total" LOR will be the sum of those two LORs.

Some totals may not appear to add up because the total is rounded after adding up the raw values.

If reported, measurement uncertainty follow the ± sign after the analytical result and is expressed as the expanded uncertainty calculated using a coverage factor of 2, providing a level of confidence of approximately 95%, unless stated otherwise in the comments section of this report.

Results reported for samples tested under test methods with codes starting with ARS-SOP, radionuclide or gross radioactivity concentrations are expressed in becquerel (Bq) per unit of mass or volume or per wipe as stated on the report. Becquerel is the SI unit for activity and equals one nuclear transformation per second.

Note that in terms of units of radioactivity:

- a. 1 Bq is equivalent to 27 pCi
- b. 37 MBq is equivalent to 1 mCi

Solid samples expressed on a dry weight basis.

For results reported for samples tested under test methods with codes starting with ARS-SOP, less than (<) values indicate the detection limit for each radionuclide or parameter for the measurement system used. The respective detection limits have been calculated in accordance with ISO 11929.

The QC and MU criteria are subject to internal review according to the SGS QAQC plan and may be provided on request or alternatively can be found here: <a href="https://www.sgs.com.au/en-gb/environment-health-and-safety">www.sgs.com.au/en-gb/environment-health-and-safety</a>.

This document is issued by the Company under its General Conditions of Service accessible at <a href="www.sgs.com/en/Terms-and-Conditions.aspx">www.sgs.com/en/Terms-and-Conditions.aspx</a>.

Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

Any holder of this document is advised that information contained hereon reflects the Company's findings at the time of its intervention only and within the limits of Client's instructions, if any. The Company's sole responsibility is to its Client only. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

This report must not be reproduced, except in full.

16-March-2021 Page 7 of 7